



MAY 2016

How To Beat Post Workout Soreness

ESSENTIAL PHYSIOTHERAPY

Most of us are familiar with the post-exercise soreness that comes after a particularly strenuous exercise session. This pain, which is usually accompanied by stiffness and weakness, is often referred to as Delayed Onset Muscle Soreness, abbreviated to DOMS.

It is unclear exactly what the purpose of this pain is, as it is most intense 24-48 hours following an intense exercise session. However, we do know that muscles experiencing DOMS show signs of microtears throughout the tissues and the effects are worse following eccentric (controlled extension) exercises more than other types.

While DOMS is not indicative of any serious injury and has no lasting side effects, it can be very uncomfortable and is problematic for anyone who needs to perform again quickly, such as an athlete in a tournament stretching over a few days.

A person suffering from DOMS may also be at a higher risk of injury if they continue to play sport at their usual level. Understandably, many people are interested in how to avoid or reduce the effects of DOMS.

There is plenty of anecdotal evidence on how to reduce DOMS, however high-quality research is limited and there is certainly no quick fix to this problem.

When it comes to avoiding DOMS, both a warm up before exercise and a cool down afterwards are important. Many experts also recommend using a foam roller on affected muscles following exercise. And while many people, particularly long-distance athletes are fond of using anti-inflammatory medication to reduce pain after exercise, the evidence shows that its effectiveness is limited and perhaps not worth the side effects of using the medication regularly.

Other advice includes gentle exercise, such as 20 minutes on a stationary bike, and mild stretching. Massage is not generally recommended, although some people may find it helpful. Staying hydrated during sports is always important, and keeping your fluid intake adequate while experiencing soreness afterwards can also help. If you are particularly brave, ice baths have recently been shown to have mild benefits in pain reduction.



Brain Teaser

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15

Can you find the mitsake?

You have 10 fish, 5 drown and 3 come back to life. How many fish do you have left?

Connect the anatomical term with its more commonly used name

Patella	Collar Bone
Clavicle	Jaw Bone
Mandible	Kneecap

PhysioTip

Diagnosing yourself online can be difficult because many conditions actually have the same symptoms.

It pays to seek professional assessment to know exactly what is causing your pain.

Shoulder Dislocations

The shoulder is an amazing joint with incredible flexibility. It doesn't attach directly to the spine, like the hip joint; instead, it is held to the body through a complicated system of musculature and indirectly by the collarbone (clavicle) to the front of the rib cage. Many other joints in the body are extremely stable, thanks to the structure of the bones and ligaments surrounding them. However, the shoulder has so much movement that some stability is sacrificed. It is for this reason that shoulder dislocations are a relatively common occurrence.

What is a dislocation and how does it happen?

As the name suggests, a dislocated shoulder is where the head of the upper arm is moved out of its normal anatomical position to sit outside of the shoulder socket joint.

Some people have more flexible

Joints than others and will unfortunately have joints that slip out of position easily without much cause. Other people might never have a dislocated shoulder except for a traumatic injury that forces it out of position. The shoulder can dislocate in many different directions, the most common being anterior. This occurs when the arm is raised and forced backwards in a 'stop sign' position, which can occur in many situations.

What to do if this happens

The first time a shoulder dislocates is usually the most serious. If the shoulder doesn't just go back in by itself (spontaneous relocation), then someone will need to help to put it back in. This needs to be done by a professional as they must be able to assess what type of dislocation has occurred, and often an X-ray needs to be taken before the relocation happens.

A small fracture can actually occur as the shoulder is being put into place, which is why it is so important to have a professional perform the procedure.


How can physiotherapy help?

Following dislocation, your physiotherapist can advise on how to allow the best healing for the shoulder. It is important to keep the shoulder protected for a period to allow any damaged structures to heal as well as they can.

After this, a muscle-strengthening and stabilization program can begin. This is aimed at helping the muscles around the shoulder to provide further stability and prevent future dislocations.

The information in this newsletter is not a replacement for proper medical advice. Always see a medical professional for assessment of your individual condition.

Dislocated Shoulder



Answers: 1. The error is the word "mistake" 2. 10, fish can't drown. 3. Patella = Kneecap, Clavicle = Collarbone, Mandible = Jaw Bone

Mushroom and Walnut Burgers



Ingredients

500g mushrooms, chopped
1 onion, diced
2 garlic cloves, crushed
150g cooked chickpeas
40g walnuts, crushed in food processor
25g basil leaves, chopped.
1 heaped teaspoon ground flaxseed
1/2 teaspoon sea salt
ground pepper to taste

1. Preheat oven to 230c. Scatter mushrooms over two baking trays and bake for about 30 mins. Set aside in a large bowl.
2. Mix ground flaxseed with enough water to cover and set aside and allow to thicken.
3. Sauté onions over medium heat until soft, add garlic and cook for about 15 minutes. Add to mushrooms along with walnuts, thickened flaxseed and chopped basil. Briefly mix together in a food processor.
4. Return mixture to bowl and add salt and pepper. Shape the mixture into four large patties and wrap each in glad wrap. Refrigerate for at least an hour.
5. When ready to cook, heat a thin layer of oil over medium heat in a heavy frying pan and cook burgers until brown.

Serve as you like, recommended with pita bread, bean sprouts, roasted red peppers and salad.



18 Olivedale St
BIRDWOOD SA, 5234

39 Queen Street
WILLIAMSTOWN SA, 5351

For Appointments:
Call (08) 8568 5455
Text 0413597417

Or Through Our Website At
www.essentialphysio.com.au

admin@essentialphysio.com.au